



Chapter (refereed)

Pickup, Roger; Rhodes, Glenn; Hermon-Taylor, John. 2010 *Mycobacterium avium* subsp. paratuberculosis, Johne's disease and Crohn's disease. In: Ayres, Jon G.; Harrison, Roy M.; Nichols, Gordon L.; Maynard, Robert L., (eds.) *Environmental Medicine*. Hodder Arnold.

Copyright © Hodder Education 2010

This version available at <http://nora.nerc.ac.uk/4813/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the authors and/or other rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the chapter, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this chapter.

www.hoddereducation.co.uk

Contact CEH NORA team at
noraceh@ceh.ac.uk

The NERC and CEH trade marks and logos ('the Trademarks') are registered trademarks of NERC in the UK and other countries, and may not be used without the prior written consent of the Trademark owner.

***Mycobacterium avium* subsp. *paratuberculosis*, Johne's disease and Crohn's disease**

R.W. Pickup¹, G. Rhodes¹ and J. Hermon-Taylor²

¹ Centre for Ecology and Hydrology, Lake Ecosystems Group, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster, Lancashire, LA1 4AP, UK. Tel: +44(0)1524 595887 (direct), Fax: +44(0)1524 61536, E-mail Roger@ceh.ac.uk

² Division of Nutritional Sciences, School of Biomedical and Health Sciences, Franklin-Wilkins Building, Kings College London, Stamford Street, London SE1 9NH. UK. Tel: +44 (0)207 848 4552, E-mail: j.hermon@kcl.ac.uk

Mycobacterium avium* subspecies *paratuberculosis *Mycobacterium avium* subspecies *paratuberculosis* (MAP) is a member of the non-tubercular *M. avium* complex (MAC) [1]. The total genome sequence comprises 4.83 Mbp of GC rich DNA encoding a predicted 4,350 ORFs with >96% of its DNA is virtually identical in sequence and genetic organisation to that of generally non-pathogenic *M. avium*. The remainder of the genome comprising <200 ORFs is either unique to MAP or homologous to genes associated with pathogenicity in other organisms[2]. MAP has diversified into multiple bovine and ovine strains [3] with evidence of the emergence of humanised strains [4]. Some strains are almost impossible to isolate by culture. Others, particularly bovine strains require prolonged incubation [5]. Overall, these are extremely difficult organisms to study experimentally.

Johne's disease in animals Unlike other MAC, MAP has the specific ability to cause Johne's disease in many animal species including primates[1]. Johne's disease is a systemic infection whose principal clinicopathological manifestation is chronic inflammation of the intestine. The disease in animals ranges from pluribacillary to paucimicrobial like Leprosy in humans [6]). Johne's disease was first reported in 1895 in Europe and in 1908 in North America. Currently there is an extensive prevalence of subclinical MAP worldwide [7]. Infected animals secrete MAP in their milk and faeces. MAP is detected in retail pasteurised milk supplies in Britain, the Czech Republic and the USA and long term in the environment (e.g. [8]).

Crohn's disease (CD) in humans Crohn's disease is a systemic disorder whose principal clinico-pathological manifestation is chronic inflammation of the intestine. It is a 'new' disease which first emerged in Europe and North America between 1940 and 1950 with generally increasing incidence and prevalence (New Zealand (2005) 16.5/10⁵ population/year prevalence of 155.2/10⁵ ; 8). Crohn's disease has also emerged and is rising in former low incidence areas e.g. China. In Europe the incidence of the disease in adults is increasing by about 25% per decade. The Czech Republic and Melbourne show that the rise in the incidence of Crohn's disease in children in recent years has averaged 5 fold per decade (9).

MAP infection in Crohn's disease MAP is strongly associated with CD [9, 10]. MAP in paucimicrobial disease in humans is extremely difficult to detect. Optimised methods showed that 92% of people with Crohn's disease were found to be infected with MAP [11], confirmed independently in Europe and the USA [12, 13]. When appropriate methods are used most people with chronic inflammation of the intestine of the Crohn's disease type are infected with these proven multi-host chronic enteric pathogens .

1. N. B. Harris, R. G. Barletta, *Clin. Microbiol. Rev.* 14, 489 (2001). 2. C. J. Clarke, *J. Comp Pathol.* 116, 217 (1997). 3. D. M. Collins, Z. M. De, S. M. Cavaignac, *J. Clin. Microbiol.* 40, 4760 (2002). 4. M. Mura *et al.*, *Appl. Environ. Microbiol.* 72, 854 (2006). 5. E. J. Manning, M. T. Collins, *Rev. Sci. Tech.* 20, 133 (2001). 6. M. Machackova *et al.*, *Vet. Microbiol.* 101, 225 (2004). 7. J. Cheng *et al.*, *World J. Microbiol. Biotechnol.* 21, 1175 (2005). 8. R. B. Gearry *et al.* *Gastroenterology* 130(4 suppl. 2), A-622. (2006). 9. R. W. Pickup *et al.*, *Appl. Environ. Microbiol.* 72, 4067 (2006). 11. J. D. Sanderson, M. T. Moss, M. L. Tizard, J. Hermon-Taylor, *Gut* 33, 890 (1992). 12. T. J. Bull *et al.*, *J. Clin. Microbiol.* 41, 2915 (2003). 15. R. W. Pickup *et al.*, *Appl. Environ. Microbiol.* 71, 2130 (2005). 17. R. D. Berghaus, *et al.*, *J. Dairy Sci.* 89, 963 (2006).

REFERENCES 1. N. B. Harris, R. G. Barletta, *Clin. Microbiol. Rev.* 14, 489 (2001) ; 2. C. J. Clarke, *J. Comp Pathol.* 116, 217 (1997). 3. D. M. Collins, Z. M. De, S. M. Cavaignac, *J. Clin. Microbiol.* 40, 4760 (2002); 4. M. Mura *et al.*, *Appl. Environ. Microbiol.* 72, 854 (2006); 5. E. J. Manning, M. T. Collins, *Rev. Sci. Tech.* 20, 133

(2001).; **6.** M. Machackova *et al.*, *Vet. Microbiol.* 101, 225 (2004); **7.** J. Cheng *et al.*, *World J. Microbiol. Biotechnol.* 21, 1175 (2005); **8.** R. B. Gearry *et al.* *Gastroenterology* 130(4 suppl. 2), A-622. (2006); **9.** R. W. Pickup *et al.*, *Appl. Environ. Microbiol.* 72, 4067 (2006); **10.** J. Hermon-Taylor, F. El-Zaatari. IWA Publishing. WHO Pathogenic Mycobacteria in Water: A guide to Public Health Consequences, Monitoring and Management. 74-94 (2004); **11.** J. D. Sanderson, M. T. Moss, M. L. Tizard, J. Hermon-Taylor, *Gut* 33, 890 (1992); **12.** T. J. Bull *et al.*, *J. Clin. Microbiol.* 41, 2915 (2003); **13.** S. A. Naser, G. Ghobrial, C. Romero, J. F. Valentine, *Lancet* 364, 1039 (2004); **14.** J.H. Feller *et al.* *Lancet Infect Dis.* 2007; 7:607-13; **15.** I. Abubakar *et al.* *Inflamm Bowel Dis* 2007 Sep 20; Epub). **16.** M. J. Daniels, N. Ball, M. R. Hutchings, A. Greig, *Vet. J.* 161, 306 (2001); **17.** R. W. Pickup *et al.*, *Appl. Environ. Microbiol.* 71, 2130 (2005); **18.** J. D. Cirillo, *et al.*, *Infect. Immun.* 65, 3759 (1997). **19.** R. D. Berghaus, *et al.*, *J. Dairy Sci.* 89, 963 (2006); **20.** E.L.Brodie *et al.* Urban aerosols harbor diverse and dynamic bacterial populations. *PNAS* 104, 299-304 (2007); **21.** S. Berger *et al.* *Clin Vaccine Immunol.* 13, 1022-1029. (2006); **22.** D. Herthnek & G. Bolske *BMC Microbiol* 4;6:87. (2006); **23.** D. Herthnek *et al.* *J Appl Microbiol.* 100:1095-1102. . (2006); **24.** A.Nocker *et al.* *Appl. Environ. Microbiol* 73, 511-5117(2007); **25.** A. Nocker *et al.* *Journal of Microbiological Methods, Volume 70, Issue 2, August 2007, Pages 252-260*; **26.** Environment Agency Climate change impacts and water temperature. Science report SC060017/SR (July 2007).

- [1] Harris ND, Barletta RG. *Mycobacterium avium subspecies paratuberculosis* in veterinary medicine. *Clinical Microbiology Reviews.* 2001;14:489-512.
- [2] Li LL, Bannantine JP, Zhang Q, Amonsin A, May BJ, Alt D, *et al.* The complete genome sequence of *Mycobacterium avium subspecies paratuberculosis*. *Proceedings of the National Academy of Sciences of the United States of America.* 2005;102(35):12344-9.
- [3] Collins DM, De Zoete M, Cavaignac SM. *Mycobacterium avium* subsp *paratuberculosis* strains from cattle and sheep can be distinguished by a PCR test based on a novel DNA sequence difference. *Journal of Clinical Microbiology.* 2002 Dec;40(12):4760-2.
- [4] Mura M, Bull TJ, Evans H, Sidi-Boumedine K, McMinn L, Rhodes G, *et al.* Replication and long-term persistence of bovine and human strains of *Mycobacterium avium* subsp *paratuberculosis* within *Acanthamoeba polyphaga*. *Applied and Environmental Microbiology.* 2006;72(1):854-9.
- [5] Pickup RW, Rhodes G, Arnott S, Sidi-Boumedine K, Bull TJ, Weightman A, *et al.* *Mycobacterium avium* subsp *paratuberculosis* in the catchment area and water of the river Taff in South Wales, United Kingdom, and its potential relationship to clustering of Crohn's disease cases in the city of Cardiff. *Applied and Environmental Microbiology.* 2005;71(4):2130-9.
- [6] Clarke CJ. The pathology and pathogenesis of *paratuberculosis* in ruminants and other species. *Journal of Comparative Pathology.* 1997;116(3):217-61.
- [7] Manning EJ, Collins MT. *Mycobacterium avium subspecies paratuberculosis*: pathogen, pathogenicity and diagnosis. *Review of Science and Technology.* 2001;20:133-50.
- [8] Millar D, Ford J, Sanderson J, Withey S, Tizard M, Doran T, *et al.* IS900 PCR to detect *Mycobacterium paratuberculosis* in retail supplies of whole pasteurized cows' milk in England and Wales. *Applied and Environmental Microbiology.* 1996;62(9):3446-52.
- [9] Feller M, Huwiler K, Stephan R, Altpeter E, Shang A, Furrer H, *et al.* *Mycobacterium avium subspecies paratuberculosis* and Crohn's disease: a systematic review and meta-analysis. *Lancet Infectious Diseases.* 2007;7(9):607-13.
- [10] Abubakar I, Myhill DJ, Hart AR, Lake IR, Harvey I, Rhodes JM, *et al.* A case-control study of drinking water and dairy products in Crohn's disease- Further investigation of the possible role of *mycobacterium avium paratuberculosis*. *American Journal of Epidemiology.* 2007;165(7):776-83.
- [11] Bull TJ, McMinn EJ, Sidi-Boumedine K, Skull A, Durkin D, Neild P, *et al.* Detection and verification of *Mycobacterium avium* subsp *paratuberculosis* in fresh ileocolonic mucosal biopsy specimens from individuals with and without Crohn's disease. *Journal of Clinical Microbiology.* 2003;41(7):2915-23.

- [12] Sechi LA, Scanu AM, Molicotti P, Cannas S, Mura M, Dettori G, et al. Detection and isolation of *Mycobacterium avium* subspecies paratuberculosis from intestinal mucosal biopsies of patients with and without Crohn's disease in Sardinia. *American Journal of Gastroenterology*. 2005;100(7):1529-36.
- [13] Naser SA, Schwartz D, Shafran I. Isolation of *Mycobacterium avium* subsp paratuberculosis from breast milk of Crohn's disease patients. *American Journal of Gastroenterology*. 2000;95(4):1094-5.